

# Safe at School'

# Diabetes Medical Management Plan

Management Plan
school YEAR:

(Add student photo here.)

STUDENT LAST NAME:

Name of Health Care Provider/Clinic:

Email Address (non-essential communication):

FIRST NAME:

DOB:

TABLE OF CONTE	NTS	
PARENT/GUARDIAN SECTIONS	PAGE	SECTION
Demographics	1	1
Supplies/Disaster Plan/Field	1	2
Trips Self-Management Skills	2	3
Student Recognition of Highs/Lows	2	4
Glucose Monitoring at School	2	5
Parent Approval Signature	6	9
DIABETES PROVIDER SECTIONS	PAGE	SECTION
Insulin Doses at School	3	6
Dosing Table (Single Page Update)	4	6A
Correction Sliding Scale	4	6B
Long Acting Insulin Other Medications	4	6C
Other Medications	4	6D
Low Glucose Prevention	5	7
Low Glucose Management	5	8
High Glucose Management	6	9
Approval Signatures	6	9

PARENTS/GUARDIANS: Please complete pages 1 and 2 of this form and approve the final plan on page 6.

1. DEMOGRAPH	IIC IIVFOR	MATION	MANENT/GU/	ANDIA	N TO COMPLET			Dete Diami	a a de
Student First Name:	Las	t Name:	DO	3:	Student's Cell #:	Diabetes Ty	pe:	Date Diagno: Month:	sed: Year:
School Name:						School Pho	ne #:	School Fax #:	Grade:
Home Room: Scho	ool Point of C	Contact:						Con	tact Phone #
STUDENT'S SCHEDU	JLE Arrival	Time:	Dism	nissal Tir	me:				
Travels to school by		Meals Tim	es:	Р	hysical Activity:		Trav	els to:	
(check all that apply):		□ Breakfa	ast		] Gym		□н	ome 🗌 After Scl	nool Progran
☐ Foot/Bicycle		☐ AM Sna	ack		Recess		v	ia: □ Foot/Bicyd	ele
☐ Car		Lunch			] Sports			☐ Car	
□ Bus		☐ PM Sna	ack		Additional informat	ion:		☐ Student Dr	river
☐ Attends Before School Program		☐ Pre Dis	missal					☐ Bus	
Parent/Guardian #1 (ca	ontact first):	YEAR LONG	Relationship:	Pi	arent/Guardian #2:			Rela	tionship:
Cell #:	Home #:		Work #:	С	ell #:	Home #:		Work #:	
E-mail Address:				E.	-mail Address:				
Indicate preferred con	tact method:			In	dicate preferred con	tact method			
2. NECESSARY	SUPPLIE	S / DISA	STER PLANNIN	IG / E	XTENDED FIEL	D TRIPS			
1. A 3-day minimum of th					View Disaster/Emerger	ncy Planning o	etails -	refer to Safe at S	chool Guide
be provided by the paren at all times.	nt/guardian and	d accessible t	or the care of the stud	3.	Please review expiration	on dates and o	uantiti	es monthly and rep	olace items
• Insulin	Meter with	(test	Cartridge, extra	pr	ior to expiration dates				
· Syringe/Pen Needles	strips, lanc	ets, extra	Battery/Charging		In the event of a disast				
Ketone Strips     Treatment for lows	battery) - r for all Cont	•	Cord) if applicable • Additional		esignated personnel will student's location.	ii take student	s diabe	ites supplies and i	nedications
and snacks	Glucose M		supplies:	10	J. Lavin v roomioth				
<ul> <li>Glucagon</li> </ul>	(CGM) use	rs							
Antiseptic Wipes     Blood Glucose (BG)	<ul> <li>Pump Sup (Infusion Set)</li> </ul>								

Contact #:

Other:

Fax #:

Name of Health Care Provider/Clinic:

Email Address (non-essential communication):

#### **Diabetes Medical Management Plan**

DOB: STUDENT LAST NAME: FIRST NAME: 3. SELF-MANAGEMENT SKILLS (DEFINITIONS BELOW) **Full Support** Supervision Self-Care Glucose Monitoring: Meter CGM (Requires Calibration) Carbohydrate Counting Syringe Insulin Administration: Pen Pump Can Calculate Insulin Doses Glucose Management: Low Glucose  $\Box$ High Glucose Self-Carry Diabetes Supplies: ☐ Yes ☐ No Please specify items: Smart Phone: ☐ Yes ☐ No Device Independence: CGM Interpretation & Alarm Management Sensor Insertion Calibration Insulin Pumps Bolus ☐ Connects/Disconnects ☐ Temp Basal Adjustment ☐ Interpretation & Alarm Management ☐ Site Insertion ☐ Cartridge Change Full Support: All care performed by school nurse and trained staff (as permitted by state law). Supervision: Trained staff to assist & supervise. Guide & encourage independence. Self-Care: Manages diabetes independently. Support is provided upon request and as needed. 4. STUDENT RECOGNITION OF HIGH OR LOW GLUCOSE SYMPTOMS (CHECK ALL THAT APPLY) **Symptoms of High:** ☐ Thirsty ☐ Frequent Urination ☐ Fatigued/Tired/Drowsy ☐ Headache ☐ Blurred Vision ☐ Warm/Dry/Flushed Skin ☐ Abdominal Discomfort ☐ Nausea/Vomiting ☐ Fruity Breath ☐ Unaware ☐ Other: Symptoms of Low: None ☐ Hungry ☐ Shaky ☐ Pale ☐ Sweaty ☐ Tired/Sleepy ☐ Tearful/Crying ☐ Dizzy Irritable ☐ Unable to Concentrate ☐ Confusion ☐ Personality Changes ☐ Other: Has student lost consciousness, experienced a seizure or required Glucagon: ☐ Yes ☐ No If yes, date of last event: Has student been admitted for DKA after diagnosis: ☐ Yes ☐ No If yes, date of last event: 5. GLUCOSE MONITORING AT SCHOOL **Monitor Glucose:** ☐ Before Meals ☐ With Physical Complaints/Illness (include ketone testing) ☐ High or Low Glucose Symptoms ☐ Before Exams ☐ Before Physical Activity ☐ After Physical Activity ☐ Before Leaving School ☐ Other: **CONTINUOUS GLUCOSE MONITORING (CGM)** Please: Permit student access to viewing device at all times (Specify Brand & Model: Permit access to School Wi-Fi for sensor data collection and data Specify Viewing Equipment: Device Reader ☐ Smart Phone ☐ Insulin Pump ☐ Tablet or iPod ☐ Smart Watch Do not discard transmitter if sensor falls CGM is remotely monitored by parent/quardian. Perform finger stick if: Document individualized communication plan in Section 504 Glucose reading is below mg/dL or above mq/dL or other plan to minimize interruptions for the student. If CGM is still reading below mg/dL (DEFAULT 70 mg/dL) ☐ May use CGM for monitoring/treatment/insulin dosing unless 15 minutes following low treatment symptoms do not match reading. CGM sensor is dislodged or sensor reading is unavailable. **CGM Alarms:** (see CGM addenda for more information) Low alarm mg/dL Sensor readings are inconsistent or in the presence of alerts/alarms Dexcom does not have both a number and arrow present High alarm mg/dL if applicable Libre displays Check Blood Glucose Symbol Using Medtronic system with Guardian sensor Notify parent/guardian if glucose is: ☐ Section 1-5 completed by Parent/Guardian below mg/dL (<55 mg/dL DEFAULT) above mg/dL (>300 mg/d DEFAULT)

Contact #:

Other:

Fax #:

STUDENT LA	AST NAME:	FIRST NAME:	DOB:
6. INSULI	N DOSES AT SCHOOL - HEALTHO	CARE PROVIDER TO COMPLE	ETE
Insulin Admi ☐ Syringe ☐ i-Port ☐ Other	nistered Via: ☐ Insulin Pen (☐ Whole Units ☐ Half Units) ☐ Smart Pen	FDA-approved device  Insulin Pump is using DIY Loop	odel: ) ed Insulin Delivery (automatic dosing) using an ing Technology (child/parent manages device with all other diabetes management)
	o be determined by Bolus Calculator in insuli evice failure (provide insulin via injection usin		oderate or large ketones are present or in the
Insulin Delive	nistration Guidelines ry Timing: Pre-meal insulin delivery is import demonstrate unpredictable eating patterns o		ol. Late or partial doses are used with arbohydrates when student does not complete
☐ After Mea	eal (DEFAULT) If as soon as possible and within 30 minutes avoid snacking hours (DEFAULT 2 hours)	ours) before and after meals	
Partial Dose	Prior to Meal: (preferred for unpredictable e	eating patterns using insulin pump the	erapy)
☐ Follow me	meal dose using grams of carbohydr al with remainder of grams of carbohydrates nce to Prior to Meal when student demonstra		I hybrid pump therapy)
For Injection	s, Calculate Insulin Dose To The Nearest:		
	round down for $< 0.25$ or $< 0.75$ and round up t (round down for $< 0.5$ and round up for $\ge 0$ .		
☐ Check for	al Insulin Orders: KETONES before administering insulin dose implains of physical symptoms. Refer to sect		00 mg/dL or >250 mg/dL on insulin pump) or if ment information.
	uardians are authorized to adjust insulin dose	+/- units	
☐ Insulin			
☐ Insulin			
	to Carb Ratio +/- grams/units		
□ Insulin	Factor +/- mg/dL/unit		
Additional gu	idance on parent adjustments:		

Safe at School

Connected for Life STUDENT LAS	Parl Comment of the Comment		FIRS	ST NAME:			D	OB:
6A. DOSIN	G TABLE—HEALTHO	ARE PROVIDI	ER TO CC	)MPLETE –	SINGLE	E PAGE UPDAT	E ORDE	R FORM
nsulin: (admin Rapid Actir Ultra Rapid	istered for food and/or corn ng Insulin: ☐ Humalog/Adr I <b>Acting Insulin:</b> ☐ Fiasp (/ in: ☐ Humulin R ☐ Novo	rection) melog (Lispro), No Aspart) 🔲 Lyumj	ovolog (Aspa	ırt), Apidra (Glı	ulisine) [			
Meal & Times	Food Dos	9		Glucose C Use Formula	orrection		□ PE/A	Activity Day Do
Select if dosing is required for meal	Carbohydrate Ratio: Total Grams of Carbohydrate divided by Carbohydrate Ra = Carbohydrate Dose	_	Glucose) di	re-Meal Glucos vided by <b>Corre</b> Correction dos DEFAULT 3 hou	ction Fact se every	minus <b>Target</b> or = Correction Dose hours as	☐ Total	ohydrate Dose Dose dose instruction
☐ Breakfast	Breakfast Carb Ratio = g/unit	Breakfast units	Correc	Glucose is: tion Factor is:		ng/dL & ng/dL/unit	Carb Ra Subtra Subtra	act %
AM Snack	AM Snack Carb Ratio = g/unit	AM Snack units	☐ Target	Glucose is: tion Factor is:		ng/dL & ng/dL/unit	Carb Ra	9
Lunch	Lunch Carb Ratio = g/unit	Lunch	☐ Target	Glucose is: tion Factor is:		ng/dL & ng/dL/unit	Carb Ra Subtra Subtra	atio g/u
☐ PM Snack	PM Snack Carb Ratio = g/unit	(CORRES	☐ Target	Glucose is:		ng/dL & ng/dL/unit	Carb Ra	act %
□ Dinner	□ No Carb Dose □ No Insu  Dinner  Carb Ratio = g/unit	Dinner	☐ Target	Glucose is: tion Factor is:		ng/dL & ng/dL/unit	Carb Ra Subtra Subtra	atio g/u
B. CORRE	ECTION SLIDING SC	ALE						
Meals Only to to to	mg/dL = unit  mg/dL = unit  mg/dL = unit  mg/dL = unit	s to	mg	i /dL = /dL = /dL =	units units units	to to	mg/dL = mg/dL = mg/dL =	units units units
	ACTING INSULIN				TE WEST		IIIg/uz =	W W W
□ Le	ntus, Basaglar, Toujeo (Glargin vemir (Detemir) esiba (Degludec) her	e)	units	☐ Daily Dose ☐ Overnight ☐ Disaster/E	Field Trip D			Subcutaneously
	R MEDICATIONS etformin		units	☐ Daily Dose ☐ Overnight ☐ Disaster/E	Field Trip D	Dose Dose	Bive 2 alik	Route
	uired here if sending page dosing update.	Diabetes Provi	ider Signatur				D	ate:

Name of Health Care Provider/Clinic:

Contact #:

Fax #:

Email Address (non-essential communication):

Other:

7 LOW CLUCOSE PREVENTION (H)	VPOCI VCEMIA)	
STUDENT LAST NAME:	FIRST NAME:	DOB:
Connected for Life		

7. LOW GLUCOSE PREVENTION (HYPOGLYCEMIA)
Allow Early Interventions
Allow Mini-Dosing of carbohydrate (i.e.,1-2 glucose tablets) when low glucose is predicted, sensor readings are dropping (down arrow) at mg/dL (DEFAULT 80 mg/dL or 120 mg/dL prior to exercise) or with symptoms.
☐ Allow student to carry and consume snacks ☐ School staff to administer
☐ Allow Trained Staff/Parent/Guardian to adjust mini dosing and snacking amounts (DEFAULT)
Insulin Management (Insulin Pumps)
Temporary Basal Rate Initiate pre-programmed rate as indicated below to avoid or treat hypoglycemia.
☐ Pre-programmed Temporary Basal Rate Named (OmniPod)
☐ Temp Target (Medtronic) ☐ Exercise Activity Setting (Tandem)
Start: minutes prior to exercise for minutes duration (DEFAULT 1 hour prior, during, and 2 hours following exercise).
Initiated by: Student Trained School Staff School Nurse
May disconnect and suspend insulin pump up to minutes (DEFAULT 60 minutes) to avoid hypoglycemia, personal injury with certain physical activities or damage to the device (keep in a cool and clean location away from direct sunlight).
Exercise (Exercise is a very important part of diabetes management and should always be encouraged and facilitated).
Exercise Glucose Monitoring
prior to exercise every 30 minutes during extended exercise following exercise with symptoms
Delay exercise if glucose is < mg/dL (120 mg/dL DEFAULT)
Pre-Exercise Routine
☐ Fixed Snack: Provide grams of carbohydrate prior to physical activity if glucose < mg/dL
☐ Added Carbs: If glucose is < mg/dL (120 DEFAULT) give grams of carbohydrates (15 DEFAULT)
□ TEMPORARY BASAL RATE as indicated above
Encourage and provide access to water for hydration, carbohydrates to treat/prevent hypoglycemia, and bathroom privileges during
physical activity
8. LOW GLUCOSE MANAGEMENT (HYPOGLYCEMIA)
Low Glucose below mg/dL (below 70 mg/dL DEFAULT) or below mg/dL before/during exercise ( DEFAULT is < 120 mg/dl).
If student is awake and able to swallow give grams of fast acting carbohydrate (DEFAULT 15 grams). Examples include 4 ounces
of juice or regular soda, 4 glucose tabs, 1 small tube glucose gel.  ☐ School nurse/parent may change amount given
2. Check blood glucose every 15 minutes and re-treat until glucose > mg/dL (DEFAULT is 80 mg/dL or 120 mg/dL before exercise).
SEVERE LOW GLUCOSE (unconscious, seizure, or unable to swallow)  Administer Glucagon, position student on their side and monitor for vomiting, call 911 and notify parent/guardian. If BG meter is available, confirm hypoglycemia via BG fingerstick. Do not delay treatment if meter is not immediately available. If wearing an insulin pump, place pump in suspend/stop mode or disconnect tubing from infusion site. Keep pump with student.
☐ Glucagon Emergency Kit by IM injection ☐ Gvoke by SC injection ☐ Auto-Injection, Gvoke HypoPen Dose: ☐ 0.5 mg or ☐ 1.0 mg
☐ Zegalogue (dasiglucagon) 0.6 mg SC by Auto-Injector ☐ Zegalogue (dasiglucagon) 0.6 mg SC by Pre-Filled Syringe
☐ Baqsimi Nasal Glucagon 3 mg

STUDENT LAST NAME:	FIRST NAME:	DOB:

#### 9. HIGH GLUCOSE MANAGEMENT (HYPERGLYCEMIA)

Management of High Glucose over mg/dL (Default is 300 mg/dL OR 250 mg/dl if on an insulin pump).

- 1. Provide and encourage consumption of water or sugar-free fluids. Give 4-8 ounces of water every 30 minutes. May consume fluids in classroom. Allow frequent bathroom privileges.
- 2. Check for Ketones (before giving insulin correction)
  - a. If Trace or Small Urine Ketones (0.1 0.5 mmol/L if measured in blood)
    - · Consider insulin correction dose. Refer to the "Correction Dose" Section 6.A-B. for designated times correction insulin may be given.
    - · Can return to class and PE unless symptomatic
    - · Recheck glucose and ketones in 2 hours
  - b. If Moderate or Large Urine Ketones (0.6 1.4 mmol/L or >1.5 mmol/L blood ketones). This may be serious and requires action.

☐ Send student's diabetes log to Health Care Provider (include details): If pre-meal blood glucose is below 70 mg/dL or above 240 mg/dL.

- · Contact parents/guardian or, if unavailable, healthcare provider
- Administer correction dose via injection. If using Automated Insulin Delivery contact parent/provider about turning off automatic pump features. Refer to the "Blood Glucose Correction Dose" Section 6.A-B
- · If using insulin pump change infusion site/cartridge or use injections until dismissal.
- · No physical activity until ketones have cleared

more than 3 times per week or you have any other concerns.

- · Report nausea, vomiting, and abdominal pain to parent/guardian to take student home.
- · Call 911 if changes in mental status and labored breathing are present and notify parents/guardians.

SIGNATURES			
This Diabetes Medical Management Plan ha	as been approved	d by:	
Student's Physician/Health Care Provider:	Date:		
I, (parent/quardian)	give permissi	on to the school nurse or another qualified hea	alth care professional o
	g.ve permiser	to perform and carry out the	
trained diabetes personnel of (school)			
	Plan. I also conse	ent to the release of the information contained	in this Diabetes Medic
outlined in this Diabetes Medical Management Management Plan to all school staff members	and other adults	who have responsibility for my child and who r	may need to know
trained diabetes personnel of (school) outlined in this Diabetes Medical Management Management Plan to all school staff members this information to maintain my child's health a	and other adults and safety. I also g	who have responsibility for my child and who r live permission to the school nurse or another	may need to know
outlined in this Diabetes Medical Management Management Plan to all school staff members	and other adults and safety. I also g	who have responsibility for my child and who r live permission to the school nurse or another	may need to know
outlined in this Diabetes Medical Management Management Plan to all school staff members this information to maintain my child's health a	and other adults and safety. I also g	who have responsibility for my child and who r live permission to the school nurse or another	may need to know

Name of Health Care Provider/Clinic:

Contact #:

Fax #:

Email Address (non-essential communication):

Other: